



Institutional Members: CEPR, NBER and Università Bocconi

WORKING PAPER SERIES

A characterization of probabilities with full support in metric spaces, and Laplace's method

Simone Cerreia-Vioglio, Fabio Maccheroni, Massimo Marinacci

Working Paper n. 620

This Version: February 21, 2018

IGIER – Università Bocconi, Via Guglielmo Röntgen 1, 20136 Milano –Italy
<http://www.igier.unibocconi.it>

The opinions expressed in the working papers are those of the authors alone, and not those of the Institute, which takes non institutional policy position, nor those of CEPR, NBER or Università Bocconi.

A characterization of probabilities with full support in metric spaces, and Laplace's method

Simone Cerreia-Vioglio, Fabio Maccheroni, Massimo Marinacci

Università Bocconi and IGIER

February 21, 2018

Abstract

We show that a probability measure on a metric space X has full support if and only if the set of all probability measures that are absolutely continuous with respect to it is dense in $\mathcal{P}(X)$. We illustrate the result through a general version of Laplace's method, which in turn leads to a general stochastic convergence result to global maxima.

1 Introduction

A probability measure λ on a finite space X has full support if and only if the set $\mathcal{P}_\lambda(X)$ of all probability measures that are absolutely continuous with respect to λ *coincides with* the set $\mathcal{P}(X)$ of all probability measures. Here we show that, on a metric space X , a probability measure λ has full support if and only if $\mathcal{P}_\lambda(X)$ *is dense in* $\mathcal{P}(X)$.

Since the assumption of full support is equivalent to strict positivity on nonempty open sets, our result sheds light on the notion of strict positivity of a probability measure in the infinite case. In a functional analysis perspective, it can be regarded as a characterization of strictly positive continuous linear functionals in the dual pair $\langle C_b(X), ca(X) \rangle$.

To illustrate this result, we prove a general version of Laplace's method. Specifically, if λ is a full support measure on the compact metrizable set K and $u \in C(K)$, then

$$v_n = \frac{1}{n} \log \int_K e^{nu(x)} d\lambda(x) \rightarrow v = \max_{x \in K} u(x)$$

By variational methods we show that, when the maximizer x^u of u on K is unique, to the sequence $\{v_n\}$ corresponds a sequence $\{\mu_n\}$ of measures on K that, eventually, concentrates on x^u . Moreover, if K is contained in a reflexive and separable Banach space, the sequence of the barycenters of μ_n weakly converges to x^u .

2 Setup and preliminaries

Let X be a topological space. We denote by $C(X)$ (resp., $C_b(X)$) the vector space of all continuous (resp., continuous and bounded) functions $f : X \rightarrow \mathbb{R}$, by $\mathcal{B}(X)$ the Borel sigma-algebra of X , and by $\mathcal{P}(X)$ the set of all Borel probability measures on $\mathcal{B}(X)$ with the topology $\sigma(\mathcal{P}(X), C_b(X))$ of weak convergence.

Given any $\lambda \in \mathcal{P}(X)$, we denote by $\mathcal{P}_\lambda(X)$ (resp., $\mathcal{P}_\lambda^*(X)$) the collection of all $\mu \in \mathcal{P}(X)$ that are absolutely continuous with respect to λ (resp., that have continuous and bounded density with respect to λ), and by $\ell_\lambda : C_b(X) \rightarrow \mathbb{R}$ the positive linear functional $\ell_\lambda(f) = \int_X f d\lambda$.

Definition 1 *The support of $\lambda \in \mathcal{P}(X)$, denoted by $\text{supp } \lambda$, is (if it exists) a closed subset of X with λ -null complement and such that $\lambda(G) > 0$ for all open subsets G of X having nonempty intersection with it.*

The probability measure λ has *full support* if $\text{supp } \lambda = X$, that is, $\lambda(G) > 0$ for all nonempty open subsets G of X .

If X is the dual of a separable normed space (for example, a reflexive and separable Banach space), we endow it with the weak* topology and consider the Borel sigma-algebra generated by this topology. With this topology, compact sets are metrizable and their closed and convex hulls are compact too.¹ The next basic result is a slight modification of Proposition 1.1 of Phelps (2001).

Proposition 1 *If $\mu \in \mathcal{P}(X)$ has bounded support, then there exists a unique element $m \in X$ such that*

$$\langle \phi, m \rangle = \int_X \langle \phi, x \rangle d\mu(x) \tag{1}$$

for all linear and continuous functionals $\phi : X \rightarrow \mathbb{R}$.

The element m , called *barycenter* of μ , belongs to the closed and convex hull of $\text{supp } \mu$. When X is \mathbb{R}^n , the barycenter of a Borel probability measure μ on \mathbb{R}^n that has bounded support is easily seen to be the vector $m = \int_X x d\mu(x)$.

3 Main result

We state and prove our main result. The equivalence between points (i) and (iv), i.e., between the strict positivity of λ and ℓ_λ , is essentially known and reported here for completeness and perspective.

¹Because of the Alaoglu's Theorem and of Theorem 6.30 of Aliprantis and Border, 2006 (henceforth, AB).

Theorem 1 *Let X be a metric space. The following conditions are equivalent for $\lambda \in \mathcal{P}(X)$:*

- (i) λ has full support X ;
- (ii) $\text{cl}(\mathcal{P}_\lambda^*(X)) = \mathcal{P}(X)$;
- (iii) $\text{cl}(\mathcal{P}_\lambda(X)) = \mathcal{P}(X)$;
- (iv) ℓ_λ is strictly positive, i.e., $\int_X f d\lambda > 0$ for all $0 \neq f \in C_b^+(X)$.

Proof If X is a singleton, the statement is trivial. Let us assume that X contains more than one point.

(i) implies (ii). We first show that $\delta_{\bar{x}} \in \text{cl}(\mathcal{P}_\lambda^*(X))$ for all $\bar{x} \in X$. Let $\bar{x} \in X$, and, for each $n \in \mathbb{N}$, consider the sets B_n and C_n defined by

$$B_n = \left\{ x \in X : d(x, \bar{x}) \leq \frac{1}{n} \right\} \text{ and } C_n = \left\{ x \in X : d(x, \bar{x}) \geq \frac{2}{n} \right\}.$$

Both sets are closed and clearly $B_n \cap C_n = \emptyset$. If n is large enough, say for all $n \geq \bar{n}$, both sets are nonempty because there exists $x \neq \bar{x}$ in X . By the Urysohn Lemma (e.g., [1, Theorem 2.46]), it follows that for each $n \geq \bar{n}$ there exists $\varphi_n \in C_b(X)$ such that $\varphi_n(X) \subseteq [0, 1]$, $\varphi_n(B_n) = 1$, and $\varphi_n(C_n) = 0$. Since $\bar{x} \in \text{supp } \lambda$ and $\varphi_n(\bar{x}) = 1$,² it follows that

$$k_n = \int_X \varphi_n d\lambda > 0 \quad \forall n \geq \bar{n}.$$

Now, for each $n \geq \bar{n}$, set $\psi_n = \varphi_n/k_n$ and define the measure $\lambda_n : \mathcal{B} \rightarrow \mathbb{R}$ by $\lambda_n(B) = \int_B \psi_n d\lambda$. Notice that $\lambda_n \in \mathcal{P}_\lambda^*(X)$ because $\psi_n \in C_b(X)$.

We next show that $\lambda_n \rightarrow \delta_{\bar{x}}$. Define $S_n = \{x \in X : d(x, \bar{x}) \leq 2/n\}$ for all $n \geq \bar{n}$. Notice that $S_n^c \subseteq C_n$ so that $1 = \int_{S_n} \psi_n d\lambda + \int_{S_n^c} \psi_n d\lambda = \int_{S_n} \psi_n d\lambda = \lambda_n(S_n)$ for all $n \geq \bar{n}$. Consider an open subset G of X . We have two cases:

1. $\bar{x} \notin G$. It follows that $\liminf \lambda_n(G) \geq 0 = \delta_{\bar{x}}(G)$.
2. $\bar{x} \in G$. For $n \geq \bar{n}$ large enough, say $n \geq \bar{m}$, we have that $S_n \subseteq G$. Then, for all $n \geq \bar{m}$, $\lambda_n(G) \geq \lambda_n(S_n) \geq 1$, yielding that $\liminf \lambda_n(G) \geq 1 = \delta_{\bar{x}}(G)$.

In both cases, $\liminf \lambda_n(G) \geq \delta_{\bar{x}}(G)$ holds. Since G was an arbitrarily chosen open subset of X , by the Portmanteau Theorem (e.g., [1, Theorem 15.3]) it follows that $\lambda_n \rightarrow \delta_{\bar{x}}$.

Since \bar{x} was arbitrarily chosen in X , we have that $\{\delta_x\}_{x \in X} \subseteq \text{cl}(\mathcal{P}_\lambda^*(X))$. Since $\mathcal{P}_\lambda^*(X)$ is convex, then $\text{cl}(\mathcal{P}_\lambda^*(X))$ is closed and convex, it follows that $\text{cl}(\mathcal{P}_\lambda^*(X)) \supseteq \text{cl}(\text{co}(\{\delta_x\}_{x \in X}))$. But

²E.g., [1, Lemma 12.16].

$\text{co}(\{\delta_x\}_{x \in X})$ is dense in $\mathcal{P}(X)$ (e.g., [1, Theorem 15.10]), we conclude that $\mathcal{P}(X) \supseteq \text{cl}(\mathcal{P}_\lambda^*(X)) \supseteq \text{cl}(\text{co}(\{\delta_x\}_{x \in X})) = \mathcal{P}(X)$.

(ii) implies (iii). This follows from $\mathcal{P}_\lambda^*(X) \subseteq \mathcal{P}_\lambda(X)$.

(iii) implies (iv). By contradiction, assume that $\text{cl}(\mathcal{P}_\lambda(X)) = \mathcal{P}(X)$ and ℓ_λ is not strictly positive. In this case, there exists $g \in C_b^+(X) \setminus \{0\}$ such that $\int_X g d\lambda = 0$. Consider the open set $G = \{x \in X : g(x) > 0\} \neq \emptyset$. Since $\int_X g d\lambda = 0$, then $\lambda(\{x \in X : g(x) > 0\}) = 0$, that is, $\lambda(G) = 0$. Consider $\bar{x} \in G$. Since $\text{cl}(\mathcal{P}_\lambda(X)) = \mathcal{P}(X)$, there exists a net $\{\lambda_\alpha\} \subseteq \mathcal{P}_\lambda(X)$ such that $\lambda_\alpha \rightarrow \delta_{\bar{x}}$. For each α , since λ_α is absolutely continuous with respect to λ , we have that $\lambda_\alpha(G) = 0$. Since $\lambda_\alpha \rightarrow \delta_{\bar{x}}$, by the Portmanteau Theorem, we have that $0 = \liminf \lambda_\alpha(G) \geq \delta_{\bar{x}}(G) = 1$, a contradiction.

(iv) implies (i). By contradiction, assume that ℓ_λ is strictly positive and there exists a nonempty open subset G of X with $\lambda(G) = 0$. Consider $\bar{x} \in G$. By the Urysohn Lemma, and since G^c is closed and nonempty, there exists $\varphi \in C_b(X)$ such that $\varphi(X) \subseteq [0, 1]$, $\varphi(\bar{x}) = 1$, and $\varphi(x) = 0$ for all $x \in G^c$. Since $\varphi \in C_b^+(X) \setminus \{0\}$, it follows that

$$0 < \ell_\lambda(\varphi) = \int_X \varphi d\lambda = \int_G \varphi d\lambda + \int_{G^c} \varphi d\lambda = 0,$$

a contradiction. ■

Finally, observe that the result depends only on the topology of X , so we could have used the term metrizable, rather than metric, throughout.

4 Illustration: Laplace's Method

Consider the optimization problem

$$\max_x u(x) \quad \text{sub } x \in K \tag{2}$$

where $u : X \rightarrow \mathbb{R}$ is a continuous function and K is a compact and metrizable set.

Laplace's Method is a fundamental method to find maximum values and maximizers of this general optimization problem. For this reason, it plays an important role in many applications (see, e.g., Parpas and Rustem, 2009, for an introductory overview and some relevant references). To illustrate the scope of our main result, here we establish a general abstract version of this classic method. A related result appears in Hwang (1980), though in a different setup and with an altogether different approach.

In the statement we denote by \xrightarrow{w} the $\sigma(\mathcal{P}(X), C_b(X))$ -convergence and by δ_x the Dirac probability measure concentrated on a point $x \in X$.

Theorem 2 Let X be a topological space, $u : X \rightarrow \mathbb{R}$ a continuous function, λ a Borel probability measure with compact and metrizable support K , and $\{s_n\} \subseteq (0, \infty)$ a divergent sequence. Then

$$\frac{1}{s_n} \log \int_X e^{s_n u} d\lambda \rightarrow \max_K u \quad \text{as } n \rightarrow \infty \quad (3)$$

Moreover, if u has a unique maximizer x^u in K ,³ then

$$\mu_n \xrightarrow{w} \delta_{x^u} \quad \text{as } n \rightarrow \infty \quad (4)$$

where μ_n is, for each $n \in \mathbb{N}$, defined by

$$\mu_n(B) = \frac{\int_B e^{s_n u} d\lambda}{\int_X e^{s_n u} d\lambda} \quad \forall B \in \mathcal{B}(X) \quad (5)$$

Proof First assume $K = X$, that is, X is compact and metrizable, and λ has full support. In this case, $\sigma(\mathcal{P}(X), C_b(X)) = \sigma(\mathcal{P}(X), C(X))$ is the relative weak* topology on $\mathcal{P}(X)$, and $\mathcal{P}(X)$ is compact and metrizable with respect to it (see Theorems 14.15 and 15.11 of AB). Denote

$$R(\mu||\lambda) = \begin{cases} \int_X \frac{d\mu}{d\lambda} \log \left(\frac{d\mu}{d\lambda} \right) d\mu & \text{if } \mu \ll \lambda \\ \infty & \text{else} \end{cases}$$

the relative entropy of any μ in $\mathcal{P}(X)$ with respect to λ (see Chapter 1.4 of Dupuis and Ellis, 1997, henceforth DE).

For each $n \in \mathbb{N}$, set $f_n = -s_n u$ and observe that, by Proposition 1.4.2 of DE,

$$-\log \int_X e^{-f_n} d\lambda = \min_{\mu \in \mathcal{P}(X)} \left\{ R(\mu||\lambda) + \int_X f_n d\mu \right\}$$

and the minimum of this variational formula is uniquely attained at the element μ_n of $\mathcal{P}(X)$ given by

$$\mu_n(B) = \frac{\int_B e^{-f_n(x)} d\lambda(x)}{\int_X e^{-f_n(y)} d\lambda(y)}$$

for all Borel subsets B of X . Recalling our substitution

$$\begin{aligned} -\frac{1}{s_n} \log \int_X e^{s_n u} d\lambda &= \frac{1}{s_n} \left[-\log \int_X e^{-f_n} d\lambda \right] = \frac{1}{s_n} \min_{\mu \in \mathcal{P}(X)} \left\{ R(\mu||\lambda) - s_n \int_X u d\mu \right\} \\ &= \min_{\mu \in \mathcal{P}(X)} \left\{ \frac{1}{s_n} R(\mu||\lambda) - \int_X u d\mu \right\} \end{aligned}$$

For each $n \in \mathbb{N}$, the function $F_n : \mathcal{P}(X) \rightarrow (-\infty, \infty]$ defined by

$$F_n(\mu) = \frac{1}{s_n} R(\mu||\lambda) - \int_X u d\mu \quad \forall \mu \in \mathcal{P}(X)$$

³A simple condition that ensures such uniqueness on convex sets is the strict quasi-concavity of u .

is weak* lower semicontinuous on $\mathcal{P}(X)$ (see Lemma 1.4.3 of DE and Proposition 1.9 of Dal Maso, 1993; henceforth, DM). Moreover, the sequence $\{F_n\}$ is decreasing and pointwise converges to

$$F_\infty(\mu) = \chi_{\text{dom } R(\cdot\|\lambda)}(\mu) - \int_X u d\mu \quad \forall \mu \in \mathcal{P}(X) \quad (6)$$

By Proposition 5.7 of DM, this sequence Γ -converges to the weak* lower semicontinuous envelope $\text{sc}^- F_\infty$ of F_∞ . Since $U : \mu \mapsto \int_X u d\mu$ is continuous and everywhere finite on $\mathcal{P}(X)$, by Proposition 3.7 and Example 3.4 of DM

$$\left(\bar{\text{sc}} F_\infty\right)(\mu) = \left(\bar{\text{sc}} \chi_{\text{dom } R(\cdot\|\lambda)}\right)(\mu) - \int_X u d\mu = \chi_{\text{cl}(\text{dom } R(\cdot\|\lambda))}(\mu) - \int_X u d\mu$$

For each $\mu \in \mathcal{P}_\lambda^*(X)$, $d\mu/d\lambda$ is bounded and continuous, hence there exists $k \geq 0$ such that $0 \leq d\mu/d\lambda \leq k$ and so

$$-\frac{1}{e} \leq \frac{d\mu}{d\lambda} \log\left(\frac{d\mu}{d\lambda}\right) \leq k^2 \implies R(\mu\|\lambda) < \infty \implies \mu \in \text{dom } R(\cdot\|\lambda)$$

Therefore $\mathcal{P}_\lambda^*(X) \subseteq \text{dom } R(\cdot\|\lambda)$ and so, by Theorem 1, $\mathcal{P}(X) = \text{cl}(\mathcal{P}_\lambda^*(X)) \subseteq \text{cl}(\text{dom } R(\cdot\|\lambda)) = \mathcal{P}(X)$. Summing up, F_n Γ -converges to $-\int_X u d\mu$. By Theorem 7.4 of DM, this implies

$$\lim_{n \rightarrow \infty} \min_{\mu \in \mathcal{P}(X)} \left\{ \frac{1}{s_n} R(\mu\|\lambda) - \int_X u d\mu \right\} = \min_{\mu \in \mathcal{P}(X)} \left\{ - \int_X u d\mu \right\} = - \max_{\mu \in \mathcal{P}(X)} \left\{ \int_X u d\mu \right\} = - \max_{x \in X} u(x)$$

But, for all $n \in \mathbb{N}$ we have

$$\min_{\mu \in \mathcal{P}(X)} \left\{ \frac{1}{s_n} R(\mu\|\lambda) - \int_X u d\mu \right\} = -\frac{1}{s_n} \log \int_X e^{s_n u} d\lambda$$

So, (3) holds.

Moreover, if u has a unique maximizer x^u in X , then U has δ_{x^u} as its unique maximizer. In fact, if $\mu \in \mathcal{P}(X) \setminus \{\delta_{x^u}\}$, then $\mu(X \setminus \{x^u\}) > 0$, and so

$$\begin{aligned} \int_X u d\delta_{x^u} - \int_X u d\mu &= \int_X (u(x^u) - u(x)) d\mu(x) \\ &= \int_{\{x^u\}} (u(x^u) - u(x)) d\mu(x) + \int_{X \setminus \{x^u\}} (u(x^u) - u(x)) d\mu(x) \end{aligned}$$

where the first summand is null, the second is strictly positive.⁴ Since $\mathcal{P}(X)$ is compact, the sequence F_n is equi-coercive (see Definition 7.6 of DM); in addition, it Γ -converges to $-U$ with unique minimum point δ_{x^u} in $\mathcal{P}(X)$. For each n , the probability measure μ_n is a minimizer for F_n in $\mathcal{P}(X)$. By Corollary 7.24 of DM, μ_n weak* converges to δ_{x^u} .

⁴ $\int_{B \setminus \{x^u\}} (u(x^u) - u(x)) d\gamma(x) = 0$ would imply $\gamma(\{x \in B \setminus \{x^u\} : u(x^u) - u(x) > 0\}) = 0$, a contradiction because $u(x^u) - u(x) > 0$ for all $x \in B \setminus \{x^u\}$.

In the general case, consider the compact and metrizable space K , the continuous function $w = u|_K$, and the Borel probability measure $\nu = \lambda|_K$. It is easy to show that ν has full support on K . In fact, if O is a nonempty open subset of K , there exists an open subset G of X such that $\emptyset \neq O = G \cap K = G \cap \text{supp } \lambda$; by definition of support, it follows $\lambda(G) > 0$, but then $\nu(O) = \lambda(G \cap \text{supp } \lambda) = \lambda(G \cap \text{supp } \lambda) + \lambda(G \cap (\text{supp } \lambda)^c) = \lambda(G) > 0$. The previous part of the proof implies

$$\frac{1}{s_n} \log \int_K e^{s_n w} d\nu \rightarrow \max_K w \quad \text{as } n \rightarrow \infty$$

But $s_n^{-1} \log \int_X e^{s_n u} d\lambda = s_n^{-1} \log \int_K e^{s_n w} d\nu$ for all $n \in \mathbb{N}$ and $\max_K u = \max_K w$, thus (3) holds.

Moreover, if u has a unique maximizer x^u in K , again by the previous part of the proof we can consider the sequence $\{\rho_n\}$ of probability measures defined by

$$\rho_n(L) = \frac{\int_L e^{s_n w} d\nu}{\int_K e^{s_n w} d\nu} \quad \forall n \in \mathbb{N}$$

for all Borel subsets L of K , and have that, given any $g \in C(K)$,

$$\int_K g d\rho_n \rightarrow g(x^u) \quad \text{as } n \rightarrow \infty$$

But for each $f \in C_b(X)$, $f|_K \in C(K)$ and $\int_X f d\mu_n = \int_K f|_K d\rho_n$ for all $n \in \mathbb{N}$, then the sequence $\{\mu_n\}$, defined by (5), $\sigma(\mathcal{P}(X), C_b(X))$ converges to δ_{x^u} . \blacksquare

If X is the dual of a separable normed space and is endowed with the weak* topology, then the boundedness of the support of λ is equivalent to its compactness, and – as we observed in the previous section – each μ_n has a barycenter m_n in the weak* closed and convex hull of $K = \text{supp } \lambda$. Next we show that these barycenters weak*-converge to the maximizer. Here $\xrightarrow{w^*}$ denotes weak*-convergence.

Proposition 2 *Let X be the dual of a separable normed space. Under the assumptions of Theorem 2, we have*

$$m_n \xrightarrow{w^*} x^u \quad \text{as } n \rightarrow \infty$$

where m_n is, for each $n \in \mathbb{N}$, the barycenter of μ_n .

In particular, if X is a separable and reflexive Banach space, then its weak and weak* topologies coincide and so m_n weakly converges to x^u . Clearly, the sequence of barycenters is included in K if this set is convex.

When X is \mathbb{R}^n and ℓ is a sigma-finite Borel measure, we have

$$\frac{1}{s_n} \log \frac{1}{\ell(K)} \int_K e^{s_n u(x)} d\ell(x) \rightarrow \max_K u \quad \text{as } n \rightarrow \infty \quad (7)$$

and, if x^u is the unique maximizer of u on K ,

$$m_n = \frac{\int_K e^{s_n u(x)} x d\ell(x)}{\int_K e^{s_n u(y)} d\ell(y)} \rightarrow x^u \quad (8)$$

This convergence in \mathbb{R}^n has been first established by Pincus (1968, 1970) (see Hiriart-Urruty, 1995, p. 22). The weak* convergence (4) thus substantially generalizes his results.

References

- [1] C. D. Aliprantis and K. C. Border, *Infinite Dimensional Analysis*, 3rd ed., Springer, New York, 2006.
- [2] G. Dal Maso, *An introduction to Γ -convergence*, Birkhäuser, Boston, 1993.
- [3] P. Dupuis and R. S. Ellis, *A weak convergence approach to the theory of large deviations*, John Wiley & Sons, New York, 2011.
- [4] J.-B. Hiriart-Urruty, Conditions for global optimality, in *Handbook of Global Optimization* (R. Horst and P. M. Pardalos, eds.), Springer, New York, 1995.
- [5] C.R. Hwang, Laplace's method revisited: weak convergence of probability measures, *Annals of Probability*, 8, 1177-1182, 1980.
- [6] P. Parpas and B. Rustem, Laplace method and applications to optimization problems, in *Encyclopedia of Optimization* (C. A. Floudas and P. M. Pardalos, eds.), Springer, New York, 2009.
- [7] M. Pincus, A closed form solution of certain programming problems, *Operations Research*, 16, 690-694, 1968.
- [8] M. Pincus, A Monte Carlo method for the approximate solution of certain types of constrained optimization problems, *Operations Research*, 18, 1225-1228, 1970.